

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 June 2006 (15.06.2006)

PCT

(10) International Publication Number  
**WO 2006/062883 A2**

(51) International Patent Classification:  
**A61K 31/46** (2006.01)

(21) International Application Number:  
PCT/US2005/043875

(22) International Filing Date:  
5 December 2005 (05.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/633,669 6 December 2004 (06.12.2004) US

(71) Applicant (for all designated States except US):  
**SMITHKLINE BEECHAM CORPORATION**  
[US/US]; One Franklin Plaza, P.O. Box 7929, Philadelphia, PA 19101 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LAINE, Dramane, Ibrahim** [FR/US]; 709 Swedeland Road, King Of Prussia, PA 19406 (US). **PALOVICH, Michael, R.** [US/US]; 709 Swedeland Road, King Of Prussia, PA 19406 (US).

(74) Agents: **SIMON, Soma, G.** et al.; GlaxoSmithKline, Corporate Intellectual Property UW2220, P.O. Box 1539, 709 Swedeland Road, King Of Prussia, PA 19406-0939 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MEDICAL COMBINATIONS

(57) Abstract: Muscarinic Acetylcholine Receptor Antagonists and methods of using them are provided.



WO 2006/062883 A2

**MEDICAL COMBINATIONS****FIELD OF THE INVENTION**

This invention relates to the olefinic derivatives of 8-azoniabicyclo[3.2.1] octanes, pharmaceutical compositions, and use thereof in treating muscarinic acetylcholine receptor mediated diseases of the respiratory tract and combinations thereof.

**BACKGROUND OF THE INVENTION**

Acetylcholine released from cholinergic neurons in the peripheral and central nervous systems affects many different biological processes through interaction with two major classes of acetylcholine receptors – the nicotinic and the muscarinic acetylcholine receptors. Muscarinic acetylcholine receptors (mAChRs) belong to the superfamily of G-protein coupled receptors that have seven transmembrane domains. There are five subtypes of mAChRs, termed M<sub>1</sub>-M<sub>5</sub>, and each is the product of a distinct gene. Each of these five subtypes displays unique pharmacological properties. Muscarinic acetylcholine receptors are widely distributed in vertebrate organs where they mediate many of the vital functions. Muscarinic receptors can mediate both inhibitory and excitatory actions. For example, in smooth muscle found in the airways, M<sub>3</sub> mAChRs mediate contractile responses. For review, please see Caulfield (1993 *Pharmac. Ther.* **58**:319-79).

In the lungs, mAChRs have been localized to smooth muscle in the trachea and bronchi, the submucosal glands, and the parasympathetic ganglia. Muscarinic receptor density is greatest in parasympathetic ganglia and then decreases in density from the submucosal glands to tracheal and then bronchial smooth muscle. Muscarinic receptors are nearly absent from the alveoli. For review of mAChR expression and function in the lungs, please see Fryer and Jacoby (1998 *Am J Respir Crit Care Med* **158**(5, pt 3) S 154-60).

Three subtypes of mAChRs have been identified as important in the lungs, M<sub>1</sub>, M<sub>2</sub> and M<sub>3</sub> mAChRs. The M<sub>3</sub> mAChRs, located on airway smooth muscle, mediate muscle contraction. Stimulation of M<sub>3</sub> mAChRs activates the enzyme phospholipase C via binding of the stimulatory G protein Gq/11 (Gs), leading to liberation of phosphatidyl inositol-4,5-bisphosphate, resulting in phosphorylation of contractile proteins. M<sub>3</sub> mAChRs are also found on pulmonary submucosal glands. Stimulation of this population of M<sub>3</sub> mAChRs results in mucus secretion.

M<sub>2</sub> mAChRs make up approximately 50-80% of the cholinergic receptor population on airway smooth muscles. Although the precise function is still unknown, they inhibit

catecholaminergic relaxation of airway smooth muscle via inhibition of cAMP generation. Neuronal M<sub>2</sub> mAChRs are located on postganglionic parasympathetic nerves. Under normal physiologic conditions, neuronal M<sub>2</sub> mAChRs provide tight control of acetylcholine release from parasympathetic nerves. Inhibitory M<sub>2</sub> mAChRs have also been demonstrated on sympathetic nerves in the lungs of some species. These receptors inhibit release of noradrenaline, thus decreasing sympathetic input to the lungs.

M<sub>1</sub> mAChRs are found in the pulmonary parasympathetic ganglia where they function to enhance neurotransmission. These receptors have also been localized to the peripheral lung parenchyma, however their function in the parenchyma is unknown.

Muscarinic acetylcholine receptor dysfunction in the lungs has been noted in a variety of different pathophysiological states. In particular, in asthma and chronic obstructive pulmonary disease (COPD), inflammatory conditions lead to loss of inhibitory M<sub>2</sub> muscarinic acetylcholine autoreceptor function on parasympathetic nerves supplying the pulmonary smooth muscle, causing increased acetylcholine release following vagal nerve stimulation (Fryer et al. 1999 *Life Sci* **64** (6-7) 449-55). This mAChR dysfunction results in airway hyperreactivity and hyperresponsiveness mediated by increased stimulation of M<sub>3</sub> mAChRs. Thus the identification of potent mAChR antagonists would be useful as therapeutics in these mAChR-mediated disease states.

COPD is an imprecise term that encompasses a variety of progressive health problems including chronic bronchitis, chronic bronchiolitis and emphysema, and it is a major cause of mortality and morbidity in the world. Smoking is the major risk factor for the development of COPD; nearly 50 million people in the U.S. alone smoke cigarettes, and an estimated 3,000 people take up the habit daily. As a result, COPD is expected to rank among the top five as a world-wide health burden by the year 2020. Inhaled anti-cholinergic therapy is currently considered the "gold standard" as first line therapy for COPD (Pauwels et al. 2001 *Am. J. Respir. Crit. Care Med.* **163**:1256-1276).

Despite the large body of evidence supporting the use of anti-cholinergic therapy for the treatment of airway hyperreactive diseases, relatively few anti-cholinergic compounds are available for use in the clinic for pulmonary indications. More specifically, in United States, Ipratropium Bromide (Atrovent<sup>®</sup>; and Combivent<sup>®</sup>, in combination with albuterol) is currently the only inhaled anti-cholinergic marketed for the treatment of airway hyperreactive diseases. While this compound is a potent anti-muscarinic agent, it is short acting, and thus must be administered as many as four times daily in order to provide relief for the COPD patient. In Europe and Asia, the long-acting anti-cholinergic Tiotropium Bromide (Spiriva<sup>®</sup>) was recently approved, however this product is currently not available in the United States. Thus, there

remains a need for novel compounds that are capable of causing blockade at mAChRs which are long acting and can be administered once-daily for the treatment of airway hyperreactive diseases such as asthma and COPD.

Since mAChRs are widely distributed throughout the body, the ability to apply anti-cholinergics locally and/or topically to the respiratory tract is particularly advantageous, as it would allow for lower doses of the drug to be utilized. Furthermore, the ability to design topically active drugs that have long duration of action, and in particular, are retained either at the receptor or by the lung, would allow the avoidance of unwanted side effects that may be seen with systemic anti-cholinergic use.


### SUMMARY OF THE INVENTION

This invention provides for a method of treating a muscarinic acetylcholine receptor (mAChR) mediated disease, wherein acetylcholine binds to an mAChR and which method comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

This invention also relates to a method of inhibiting the binding of acetylcholine to its receptors in a mammal in need thereof which comprises administering to aforementioned mammal an effective amount of a compound of Formula (I).

The present invention also provides for the novel compounds of Formula (I), and pharmaceutical compositions comprising a compound of Formula (I), and a pharmaceutical carrier or diluent.

The compounds according to this invention have the structure shown by Formula (I):



(I)

in which the preferred orientation of the alkyl chain attached to the tropane ring is endo.

R<sup>2</sup> and R<sup>3</sup> are, independently, selected from the group consisting of straight or branched chain lower alkyl groups having preferably from 1 to 6 carbon atoms, cycloalkyl groups having from 5 to 6 carbon atoms, cycloalkyl-alkyl having 6 to 10 carbon atoms, 2-

thienyl, 2-pyridyl, phenyl, phenyl substituted with an alkyl group having not in excess of 4 carbon atoms and phenyl substituted with an alkoxy group having not in excess of 4 carbon atoms.

X<sup>-</sup> represents an anion associated with the positive charge of the N atom. X<sup>-</sup> may be but is not limited to chloride, bromide, iodide, sulfate, benzene sulfonate, and toluene sulfonate.

Illustrative examples of this invention include

(3-*endo*)-3-(2,2-di-2-thienylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide;

(3-*endo*)-3-(2,2-diphenylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide;

(3-*endo*)-3-(2,2-diphenylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane 4-methylbenzenesulfonate;

(3-*endo*)-8,8-dimethyl-3-[2-phenyl-2-(2-thienyl)ethenyl]-8-azoniabicyclo[3.2.1]octane bromide; and

(3-*endo*)-8,8-dimethyl-3-[2-phenyl-2-(2-pyridinyl)ethenyl]-8-azoniabicyclo[3.2.1]octane bromide.

#### Combinations:

The compounds, salts and/or pharmaceutical compositions according to the present invention may also be used in combination with another therapeutically active agent, for example, a  $\beta_2$  adrenoreceptor agonist, an anti-histamine, an anti-allergic or an anti-inflammatory agent.

The present invention thus provides, in a further aspect, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with another therapeutically active agent, for example, a  $\beta_2$ -adrenoreceptor agonist, an anti-histamine, an anti-allergic, an anti-inflammatory agent or an anti-infective agent.

Examples of  $\beta_2$ -adrenoreceptor agonists include salmeterol (eg as racemate or a single enantiomer such as the R-enantiomer), salbutamol, formoterol, salmefamol, fenoterol or terbutaline, or a salt thereof (e.g. pharmaceutically acceptable salt thereof), for example the xinafoate salt of salmeterol, the sulphate salt or free base of salbutamol or the fumarate salt of formoterol. Further examples of  $\beta_2$ -adrenoreceptor agonists include, but are not limited to:

3-(4-{{6-((2*R*)-2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl)amino}hexyl}oxy}butyl)benzenesulfonamide;

3-(3-{{7-((2*R*)-2-hydroxy-2-[4-hydroxy-3-hydroxymethyl)phenyl]ethyl)-amino}heptyl)oxy}propyl)benzenesulfonamide;

4-((1*R*)-2-[(6-{2-[(2,6-dichlorobenzyl)oxy]ethoxy}hexyl)amino]-1-hydroxyethyl)-2-(hydroxymethyl)phenol

4-((1*R*)-2-[(6-{4-[3-(cyclopentylsulfonyl)phenyl]butoxy}hexyl)amino]-1-hydroxyethyl)-2-(hydroxymethyl)phenol,

5 N-[2-hydroxy-5-[(1*R*)-1-hydroxy-2-[[2-4-[(2*R*)-2-hydroxy-2-phenylethyl]amino]phenyl]ethyl]amino]ethyl]phenyl]formamide, and

N-{2-[4-(3-phenyl-4-methoxyphenyl)aminophenyl]ethyl}-2-hydroxy-2-(8-hydroxy-2(1*H*)-quinolinon-5-yl)ethylamine.

Salmeterol or a pharmaceutically acceptable salt thereof, e.g. salmeterol xinofoate, is  
10 preferably administered to humans at an inhaled dose of 25 to 50 micrograms twice per day (measured as the free base).

A combination of a compound of formula (I) or salt together with an anti-histamine is preferably for oral administration (e.g. as a combined composition such as a combined tablet), and can be for treatment and/or prophylaxis of allergic rhinitis. Examples of anti-  
15 histamines include methapyrilene, or H1 antagonists such as cetirizine, loratadine (e.g. Clarityn <sup>TM</sup>), desloratadine (e.g. Clarinex <sup>TM</sup>) or fexofenadine (e.g. Allegra <sup>TM</sup>).

Other suitable combinations include, for example, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with another anti-inflammatory agent such as an anti-inflammatory corticosteroid; or a non-steroidal anti-inflammatory drug (NSAID) such as a leukotriene antagonist (e.g. montelukast),  
20 phosphodiesterase (PDE) inhibitors (e.g. theophylline, PDE4 inhibitors or mixed PDE3/PDE4 inhibitors), an iNOS inhibitor, a tryptase inhibitor, an elastase inhibitor, a beta-2 integrin antagonist, an adenosine 2a agonist, a CCR3 antagonist, or a 5-lipoxygenase inhibitor; or an antiinfective agent (eg. an antibiotic or an antiviral). An iNOS inhibitor is preferably for oral  
25 administration. Suitable iNOS inhibitors (inducible nitric oxide synthase inhibitors) include those disclosed in WO 93/13055, WO 98/30537, WO 02/50021, WO 95/34534 and WO 99/62875. Suitable CCR3 inhibitors include those disclosed in WO 02/26722.

Of particular interest is use of the compound of formula (I) in combination with a phosphodiesterase 4 (PDE4) inhibitor or a mixed PDE3/PDE4 inhibitor. The PDE4-specific  
30 inhibitor useful in this aspect of the invention may be any compound that is known to inhibit the PDE4 enzyme or which is discovered to act as a PDE4 inhibitor, and which are only PDE4 inhibitors, not compounds which inhibit other members of the PDE family as well as PDE4. Generally it is preferred to use a PDE4 inhibitor which has an IC<sub>50</sub> ratio of about 0.1 or greater as regards the IC<sub>50</sub> for the PDE4 catalytic form which binds rolipram with a high  
35 affinity divided by the IC<sub>50</sub> for the form which binds rolipram with a low affinity. For the

purposes of this disclosure, the cAMP catalytic site which binds R and S rolipram with a low affinity is denominated the "low affinity" binding site (LPDE 4) and the other form of this catalytic site which binds rolipram with a high affinity is denominated the "high affinity" binding site (HPDE 4). This term "HPDE4" should not be confused with the term "hPDE4" which is used to denote human PDE4.

A method for determining IC<sub>50</sub>s ratios is set out in US patent 5,998,428 which is incorporated herein in full by reference as though set out herein. See also PCT application WO 00/51599 for an another description of said assay.

The preferred PDE4 inhibitors of use in this invention will be those compounds which have a salutary therapeutic ratio, i.e., compounds which preferentially inhibit cAMP catalytic activity where the enzyme is in the form that binds rolipram with a low affinity, thereby reducing the side effects which apparently are linked to inhibiting the form which binds rolipram with a high affinity. Another way to state this is that the preferred compounds will have an IC<sub>50</sub> ratio of about 0.1 or greater as regards the IC<sub>50</sub> for the PDE4 catalytic form which binds rolipram with a high affinity divided by the IC<sub>50</sub> for the form which binds rolipram with a low affinity.

A further refinement of this standard is that of one wherein the PDE4 inhibitor has an IC<sub>50</sub> ratio of about 0.1 or greater; said ratio is the ratio of the IC<sub>50</sub> value for competing with the binding of 1nM of [<sup>3</sup>H]R-rolipram to a form of PDE4 which binds rolipram with a high affinity over the IC<sub>50</sub> value for inhibiting the PDE4 catalytic activity of a form which binds rolipram with a low affinity using 1 μM[<sup>3</sup>H]-cAMP as the substrate.

Most preferred are those PDE4 inhibitors which have an IC<sub>50</sub> ratio of greater than 0.5, and particularly those compounds having a ratio of greater than 1.0. Preferred compounds are *cis* 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-carboxylic acid, 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-one and *cis*-[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol]; these are examples of compounds which bind preferentially to the low affinity binding site and which have an IC<sub>50</sub> ratio of 0.1 or greater.

Other compounds of interest include:

Compounds set out in U.S. patent 5,552,438 issued 03 September, 1996; this patent and the compounds it discloses are incorporated herein in full by reference. The compound of particular interest, which is disclosed in U.S. patent 5,552,438, is *cis*-4-cyano-4-[3-(cyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid (also known as cilomast) and its salts, esters, pro-drugs or physical forms;

AWD-12-281 from elbion (Hofgen, N. et al. 15th EFMC Int Symp Med Chem (Sept 6-10, Edinburgh) 1998, Abst P.98; CAS reference No. 247584020-9); a 9-benzyladenine derivative nominated NCS-613 (INSERM); D-4418 from Chiroscience and Schering-Plough; a benzodiazepine PDE4 inhibitor identified as CI-1018 (PD-168787) and attributed to Pfizer; a benzodioxole derivative disclosed by Kyowa Hakko in WO99/16766; K-34 from Kyowa Hakko; V-11294A from Napp (Landells, L.J. et al. Eur Resp J [Annu Cong Eur Resp Soc (Sept 19-23, Geneva) 1998] 1998, 12 (Suppl. 28): Abst P2393); roflumilast (CAS reference No 162401-32-3) and a pthalazinone (WO99/47505, the disclosure of which is hereby incorporated by reference) from Byk-Gulden; Pumafentrine, (-)-p-[(4*aR*\*,10*bS*\*)-9-ethoxy-1,2,3,4,4*a*,10*b*-hexahydro-8-methoxy-2-methylbenzo[*c*][1,6]naphthyridin-6-yl]-N,N-diisopropylbenzamide which is a mixed PDE3/PDE4 inhibitor which has been prepared and published on by Byk-Gulden, now Altana; arofylline under development by Almirall-Prodesfarma; VM554/UM565 from Vernalis; or T-440 (Tanabe Seiyaku; Fuji, K. et al. J Pharmacol Exp Ther, 1998, 284(1): 162), and T2585.

Other possible PDE-4 and mixed PDE3/PDE4 inhibitors include those listed in WO01/13953, the disclosure of which is hereby incorporated by reference.

A further combination according to the present invention comprises a compound of formula (I) or a pharmaceutically acceptable salt thereof together with an anti-inflammatory corticosteroid. Said anti-inflammatory corticosteroid is itself preferably for treatment and/or prophylaxis of asthma, COPD or allergic rhinitis. Said anti-inflammatory corticosteroid may for example be selected from fluticasone, fluticasone propionate (e.g. see US patent 4,335,121), beclomethasone, beclomethasone 17-propionate ester, beclomethasone 17,21-dipropionate ester, dexamethasone or an ester thereof, mometasone or an ester thereof, ciclesonide, budesonide, flunisolide, preferably 6 $\alpha$ ,9 $\alpha$ -difluoro-17 $\alpha$ -[(2-furanylcarbonyl)oxy]-11 $\beta$ -hydroxy-16 $\alpha$ -methyl-3-oxo-androsta-1,4-diene-17 $\beta$ -carbothioic acid *S*-fluoromethyl ester} or 6 $\alpha$ ,9 $\alpha$ -difluoro-11 $\beta$ -hydroxy-16 $\alpha$ -methyl-17 $\alpha$ -[(4-methyl-1,3-thiazole-5-carbonyl)oxy]-3-oxo-androsta-1,4-diene-17 $\beta$ -carbothioic acid *S*-fluoromethyl ester}, or a pharmaceutically acceptable salt thereof.



The anti-inflammatory corticosteroid is preferably for intranasal or inhaled administration. Fluticasone propionate is preferred and is preferably for inhaled administration to a human either (a) at a dose of 250 micrograms once per day or (b) at a dose of 50 to 250 micrograms twice per day.

5

Also provided is a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with  $\beta_2$ -adrenoreceptor agonist and an anti-inflammatory corticosteroid. Preferably this combination is for treatment and/or prophylaxis of asthma, COPD or allergic rhinitis. The  $\beta_2$ -adrenoreceptor agonist and/or the anti-inflammatory

10 corticosteroid can be as described above.

Most preferably, in this "triple" combination, the  $\beta_2$ -adrenoreceptor agonist is salmeterol or a pharmaceutically acceptable salt thereof (e.g. salmeterol xinafoate) and the anti-inflammatory corticosteroid is fluticasone propionate.

15 The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical composition and thus a pharmaceutical composition comprising a combination as defined above together with one or more pharmaceutically acceptable carriers and/or excipients represent a further aspect of the invention.

20 The individual compounds of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical composition(s). Appropriate doses of known therapeutic agents will readily be appreciated by those skilled in the art.

In one embodiment, the combination as defined herein can be for simultaneous inhaled  
25 administration and is disposed in a combination inhalation device. Such a combination inhalation device is another aspect of the invention. Such a combination inhalation device can comprise a combined pharmaceutical composition for simultaneous inhaled administration (e.g. dry powder composition), the composition comprising all the individual compounds of the combination, and the composition being incorporated into a plurality of  
30 sealed dose containers mounted longitudinally in a strip or ribbon inside the inhalation device, the containers being rupturable or peel-openable on demand; for example such inhalation device can be substantially as described in GB 2,242,134 A (DISKUS<sup>TM</sup>) and/or as described above. Alternatively, the combination inhalation device can be such that the individual compounds of the combination are administrable simultaneously but are stored  
35 separately (or wholly or partly stored separately for triple combinations), e.g. in separate

pharmaceutical compositions, for example as described in PCT/EP03/00598 filed on 22 January 2003, published as WO 03/061743 (e.g. as described in the claims thereof e.g. claim 1), incorporated herein in its entirety by reference.

- 5 The invention also provides a method of preparing a combination as defined herein, the method comprising either
- (a) preparing a separate pharmaceutical composition for administration of the individual compounds of the combination either sequentially or simultaneously, or
  - (b) preparing a combined pharmaceutical composition for administration of the
- 10 individual compounds of the combination simultaneously,
- wherein the pharmaceutical composition comprises the combination together with one or more pharmaceutically acceptable carriers and/or excipients.

The invention also provides a combination as defined herein, prepared by a method as

15 defined herein.

### **METHODS OF PREPARATION**

20

The compounds of Formula (I) may be obtained by applying synthetic procedures well known in the art as described in the patent US2800482, incorporated herein in its entirety by reference.

25

### **SYNTHETIC EXAMPLES**

The above synthetic examples in this invention are referenced to the examples described in the patent US2800482, incorporated herein in its entirety by reference.

30

### **BIOLOGICAL EXAMPLES**

The inhibitory effects of compounds at the M<sub>3</sub> mAChR of the present invention are determined by the following *in vitro* and *in vivo* functional assays:

35 **Analysis of Inhibition of Receptor Activation by Calcium Mobilization:**

Stimulation of mAChRs expressed on CHO cells were analyzed by monitoring receptor-activated calcium mobilization as previously described (H. M. Sarau *et al*, 1999. *Mol. Pharmacol.* 56, 657-663). CHO cells stably expressing M<sub>3</sub> mAChRs were plated in 96 well black wall/clear bottom plates. After 18 to 24 hours, media was aspirated and replaced with 100 µl of load media (EMEM with Earl's salts, 0.1% RIA-grade BSA (Sigma, St. Louis MO), and 4 µM Fluo-3-acetoxymethyl ester fluorescent indicator dye (Fluo-3 AM, Molecular Probes, Eugene, OR) and incubated 1 hr at 37° C. The dye-containing media was then aspirated, replaced with fresh media (without Fluo-3 AM), and cells were incubated for 10 minutes at 37° C. Cells were then washed 3 times and incubated for 10 minutes at 37° C in 100 µl of assay buffer (0.1% gelatin (Sigma), 120 mM NaCl, 4.6 mM KCl, 1 mM KH<sub>2</sub> PO<sub>4</sub>, 25 mM NaH CO<sub>3</sub>, 1.0 mM CaCl<sub>2</sub>, 1.1 mM MgCl<sub>2</sub>, 11 mM glucose, 20mM HEPES (pH 7.4)). 50 µl of compound ( $1 \times 10^{-11}$  –  $1 \times 10^{-5}$  M final in the assay) was added and the plates were incubated for 10 min. at 37° C. Plates were then placed into a fluorescent light intensity plate reader (FLIPR, Molecular Probes) where the dye loaded cells were exposed to excitation light (488 nm) from a 6 watt argon laser. Cells were activated by adding 50 µl of acetylcholine (0.1-10 nM final), prepared in buffer containing 0.1% BSA, at a rate of 50 µl/sec. Calcium mobilization, monitored as change in cytosolic calcium concentration, was measured as change in 566 nm emission intensity. The change in emission intensity is directly related to cytosolic calcium levels. The emitted fluorescence from all 96 wells is measured simultaneously using a cooled CCD camera. Data points are collected every second. This data was then plotting and analyzed using GraphPad PRISM software.

### Muscarinic Receptor Radioligand Binding Assays

Radioligand binding studies using 0.5 nM [<sup>3</sup>H]-N-methyl scopolamine (NMS) in a SPA format is used to assess binding of muscarinic antagonists to M<sub>1</sub>, M<sub>2</sub>, M<sub>3</sub>, M<sub>4</sub> and M<sub>5</sub> muscarinic acetylcholine receptors. In a 96-well plate, the SPA beads are pre-incubated with receptor-containing membrane for 30 min at 4°C. Then 50 mM HEPES and the test compound are added and incubated at room temperature (shaking) for 2 hours. The beads are then spun down and counted using a scintillation counter.

### Evaluation of potency and duration of action in isolated guinea pig trachea

Tracheae were removed from adult male Hartley guinea pigs (Charles River, Raleigh, NC; 400-600 grams) and placed into modified Krebs-Henseleit solution. Composition of the solution was (mM): NaCl 113.0, KCl 4.8, CaCl<sub>2</sub> 2.5, KH<sub>2</sub>PO<sub>4</sub> 1.2, MgSO<sub>4</sub> 1.2, NaHCO<sub>3</sub> 25.0

and dextrose 11.0. which was gassed with 95% O<sub>2</sub>: 5% CO<sub>2</sub> and maintained at 37°C. Each trachea was cleaned of adherent tissue and opened lengthwise. Epithelium was removed by gently rubbing the luminal surface with a cotton-tipped applicator. Individual strips were cut, approximately 2 cartilage rings in width, and suspended via silk suture in 10-ml water-jacketed organ baths containing Krebs-Henseleit solution and connected to Grass FT03C force-displacement transducers. Mechanical responses were recorded isometrically by MP100WS/Acknowledge data acquisition system (BIOPAC Systems, Goleta, CA, www.biopac.com) run on Apple G4 computers. The tissues were equilibrated under a resting tension of 1.5 g, determined to be optimal by length-tension evaluation, and washed with Krebs-Henseleit solution every 15 minutes for one hour. After the equilibration period pulmonary tissues were contracted with 10 uM carbachol until reaching plateau, which served as a reference contraction for data analysis. Tissues were then rinsed every 15 minutes over 1 hour until reaching baseline tone. The preparations were then left for at least 30 minutes before the start of the experiment.

Concentration-response curves were obtained by a cumulative addition of carbachol in half-log increments (Van Rossum, 1963, Arch. Int. Pharmacodyn., 143:299), initiated at 1 nM. Each concentration was left in contact with the preparation until the response plateaued before the addition of the subsequent carbachol concentration. Paired tissues were exposed to mAChR antagonist compounds or vehicle for 30 min before carbachol cumulative concentration-response curves were generated. All data is given as mean  $\pm$  standard error of the mean (s.e.m.) with *n* being the number of different animals.

For superfusion (duration of action) studies, the tissues were continuously superfused with Krebs-Henseleit solution at 2 ml/min for the duration of the experiment. Stock solutions of agonist and antagonist were infused (0.02 ml/min) via 22-gauge needle inserted into the superfusion tubing. Mechanical responses were recorded isometrically using a commercially-available data acquisition system (MP100WS/Acknowledge; BIOPAC Systems, Goleta, CA, www.biopac.com) interfaced with a Macintosh G4 computer (Apple, Cupertino, CA www.apple.com). The tissues were suspended under an optimal resting tension of 1.5 g. After a 60 min equilibration period, the tissues were contracted with carbachol (1 uM) for the duration of the experiment. Upon reaching a sustained contraction isoproterenol (10 uM) was administered to maximally relax the tissue, and this change served as a reference. Isoproterenol exposure was halted and the carbachol-induced tension allowed to recover. Muscarinic receptor antagonists infused at a single concentration per tissue until a sustained level of inhibition was attained. The compound was then removed and, once again, the carbachol-induced tension was allowed to recover.

The following parameters were determined for each concentration of antagonist, and expressed as the mean  $\pm$  S.E.M. for  $n$  individual animals. Inhibition of the carbachol-induced contraction was expressed as a percent of the reference response (isoproterenol) and the time required to reach one-half of this relaxation was measured (onset of response). The tension recovery following removal of the compound was determined as was the time required to reach one-half of the maximum tension recovery (offset of response). At 60 and 180 minutes after removal of the antagonist the remaining level of inhibition was determined and expressed as a percent of the isoproterenol reference.

Antagonist concentration-response curves were obtained by plotting the maximal relaxation data at 0, 60 and 180-min following antagonist withdrawal. Recovery, termed shift, was calculated from the ratio of the 0-min inhibition curve  $IC_{50}$  and the concentration of compound yielding a similar tension recovery at 60 and 180 minutes.

Halftimes for onset and offset of response were plotted vs. corresponding concentration and the data were fit with non-linear regression. These values were extrapolated at the  $IC_{50}$  (determined from the inhibition concentration-response curve) and designated  $Ot_{50}$  (time required, at the  $IC_{50}$  concentration, to reach half of the onset response) and  $Rt_{50}$  (time required, at the  $IC_{50}$  concentration, to reach half of the recovery response).

#### **Methacholine-induced bronchoconstriction – potency and duration of action**

Airway responsiveness to methacholine was determined in awake, unrestrained Balb C mice ( $n = 6$  each group). Barometric plethysmography was used to measure enhanced pause (Penh), a unitless measure that has been shown to correlate with the changes in airway resistance that occur during bronchial challenge with methacholine(2). Mice were pre-treated with 50  $\mu$ l of compound (0.003-10  $\mu$ g/mouse) in 50  $\mu$ l of vehicle (10% DMSO) intranasally (i.n.) and were then placed in the plethysmography chamber a given amount of time following drug administration (15 min – 96 h). For potency determination, a dose response to a given drug was performed, and all measurements were taken 15 min following i.n. drug administration. For duration of action determination, measurements were taken anywhere from 15 min to 96 hours following i.n. drug administration.

Once in the chamber, the mice were allowed to equilibrate for 10 min before taking a baseline Penh measurement for 5 minutes. Mice were then challenged with an aerosol of methacholine (10 mg/ml) for 2 minutes. Penh was recorded continuously for 7 min starting at the inception of the methacholine aerosol, and continuing for 5 minutes afterward. Data for

each mouse were analyzed and plotted by using GraphPad PRISM software. This experiment allows the determination of duration of activity of the administered compound.

The present compounds are useful for treating a variety of indications, including but not limited to respiratory-tract disorders such as chronic obstructive lung disease, chronic bronchitis, asthma, chronic respiratory obstruction, pulmonary fibrosis, pulmonary emphysema, and allergic rhinitis.

### **FORMULATION-ADMINISTRATION**

Accordingly, the present invention further provides a pharmaceutical formulation comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative (e.g., salts and esters) thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.

Hereinafter, the term "active ingredient" means a compound of formula (I), or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

Compounds of formula (I) will be administered via inhalation via the mouth or nose.

Dry powder compositions for topical delivery to the lung by inhalation may, for example, be presented in capsules and cartridges of for example gelatine, or blisters of for example laminated aluminium foil, for use in an inhaler or insufflator. Powder blend formulations generally contain a powder mix for inhalation of the compound of the invention and a suitable powder base (carrier/diluent/excipient substance) such as mono-, di- or polysaccharides (e.g., lactose or starch), organic or inorganic salts (e.g., calcium chloride, calcium phosphate or sodium chloride), polyalcohols (e.g., mannitol), or mixtures thereof, alternatively with one or more additional materials, such additives included in the blend formulation to improve chemical and/or physical stability or performance of the formulation, as discussed below, or mixtures thereof. Use of lactose is preferred. Each capsule or cartridge may generally contain between 20µg-10mg of the compound of formula (I) optionally in combination with another therapeutically active ingredient. Alternatively, the compound of the invention may be presented without excipients, or may be formed into particles comprising the compound, optionally other therapeutically active materials, and excipient materials, such as by co-precipitation or coating.

Suitably, the medicament dispenser is of a type selected from the group consisting of a reservoir dry powder inhaler (RDPI), a multi-dose dry powder inhaler (MDPI), and a metered dose inhaler (MDI).

By reservoir dry powder inhaler (RDPI) it is meant as an inhaler having a reservoir form pack suitable for comprising multiple (un-metered doses) of medicament in dry powder

form and including means for metering medicament dose from the reservoir to a delivery position. The metering means may for example comprise a metering cup or perforated plate, which is movable from a first position where the cup may be filled with medicament from the reservoir to a second position where the metered medicament dose is made available to the patient for inhalation.

By multi-dose dry powder inhaler (MDPI) is meant an inhaler suitable for dispensing medicament in dry powder form, wherein the medicament is comprised within a multi-dose pack containing (or otherwise carrying) multiple, defined doses (or parts thereof) of medicament. In a preferred aspect, the carrier has a blister pack form, but it could also, for example, comprise a capsule-based pack form or a carrier onto which medicament has been applied by any suitable process including printing, painting and vacuum occlusion.

The formulation can be pre-metered (eg as in Diskus, see GB 2242134 or Diskhaler, see GB 2178965, 2129691 and 2169265) or metered in use (eg as in Turbuhaler, see EP 69715). An example of a unit-dose device is Rotahaler (see GB 2064336). The Diskus inhalation device comprises an elongate strip formed from a base sheet having a plurality of recesses spaced along its length and a lid sheet hermetically but peelably sealed thereto to define a plurality of containers, each container having therein an inhalable formulation containing a compound of formula (I) preferably combined with lactose. Preferably, the strip is sufficiently flexible to be wound into a roll. The lid sheet and base sheet will preferably have leading end portions which are not sealed to one another and at least one of the said leading end portions is constructed to be attached to a winding means. Also, preferably the hermetic seal between the base and lid sheets extends over their whole width. The lid sheet may preferably be peeled from the base sheet in a longitudinal direction from a first end of the said base sheet.

In one aspect, the multi-dose pack is a blister pack comprising multiple blisters for containment of medicament in dry powder form. The blisters are typically arranged in regular fashion for ease of release of medicament therefrom.

In one aspect, the multi-dose blister pack comprises plural blisters arranged in generally circular fashion on a disk-form blister pack. In another aspect, the multi-dose blister pack is elongate in form, for example comprising a strip or a tape.

Preferably, the multi-dose blister pack is defined between two members peelably secured to one another. US Patents Nos. 5,860,419, 5,873,360 and 5,590,645 describe medicament packs of this general type. In this aspect, the device is usually provided with an opening station comprising peeling means for peeling the members apart to access each medicament dose. Suitably, the device is adapted for use where the peelable members are elongate sheets which define a plurality of medicament containers spaced along the length

thereof, the device being provided with indexing means for indexing each container in turn. More preferably, the device is adapted for use where one of the sheets is a base sheet having a plurality of pockets therein, and the other of the sheets is a lid sheet, each pocket and the adjacent part of the lid sheet defining a respective one of the containers, the device comprising driving means for pulling the lid sheet and base sheet apart at the opening station.

By metered dose inhaler (MDI) it is meant a medicament dispenser suitable for dispensing medicament in aerosol form, wherein the medicament is comprised in an aerosol container suitable for containing a propellant-based aerosol medicament formulation. The aerosol container is typically provided with a metering valve, for example a slide valve, for release of the aerosol form medicament formulation to the patient. The aerosol container is generally designed to deliver a predetermined dose of medicament upon each actuation by means of the valve, which can be opened either by depressing the valve while the container is held stationary or by depressing the container while the valve is held stationary.

Spray compositions for topical delivery to the lung by inhalation may for example be formulated as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, such as a metered dose inhaler, with the use of a suitable liquefied propellant. Aerosol compositions suitable for inhalation can be either a suspension or a solution and generally contain the compound of formula (I) optionally in combination with another therapeutically active ingredient and a suitable propellant such as a fluorocarbon or hydrogen-containing chlorofluorocarbon or mixtures thereof, particularly hydrofluoroalkanes, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetra-fluoroethane, especially 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoro-n-propane or a mixture thereof. Carbon dioxide or other suitable gas may also be used as propellant. The aerosol composition may be excipient free or may optionally contain additional formulation excipients well known in the art such as surfactants eg oleic acid or lecithin and cosolvents eg ethanol. Pressurised formulations will generally be retained in a canister (eg an aluminium canister) closed with a valve (eg a metering valve) and fitted into an actuator provided with a mouthpiece.

Medicaments for administration by inhalation desirably have a controlled particle size. The optimum aerodynamic particle size for inhalation into the bronchial system for localized delivery to the lung is usually 1-10 $\mu$ m, preferably 2-5 $\mu$ m. The optimum aerodynamic particle size for inhalation into the alveolar region for achieving systemic delivery to the lung is approximately .5-3  $\mu$ m, preferably 1-3  $\mu$ m. Particles having an aerodynamic size above 20 $\mu$ m are generally too large when inhaled to reach the small airways. Average aerodynamic



particle size of a formulation may be measured by, for example cascade impaction. Average geometric particle size may be measured, for example by laser diffraction, optical means.

To achieve a desired particle size, the particles of the active ingredient as produced may be size reduced by conventional means eg by controlled crystallization, micronisation or nanomilling. The desired fraction may be separated out by air classification. Alternatively, particles of the desired size may be directly produced, for example by spray drying, controlling the spray drying parameters to generate particles of the desired size range. Preferably, the particles will be crystalline, although amorphous material may also be employed where desirable. When an excipient such as lactose is employed, generally, the particle size of the excipient will be much greater than the inhaled medicament within the present invention, such that the "coarse" carrier is non-respirable. When the excipient is lactose it will typically be present as milled lactose, wherein not more than 85% of lactose particles will have a MMD of 60-90µm and not less than 15% will have a MMD of less than 15µm. Additive materials in a dry powder blend in addition to the carrier may be either respirable, i.e., aerodynamically less than 10 microns, or non-respirable, i.e., aerodynamically greater than 10 microns.

Suitable additive materials which may be employed include amino acids, such as leucine; water soluble or water insoluble, natural or synthetic surfactants, such as lecithin (e.g., soya lecithin) and solid state fatty acids (e.g., lauric, palmitic, and stearic acids) and derivatives thereof (such as salts and esters); phosphatidylcholines; sugar esters. Additive materials may also include colorants, taste masking agents (e.g., saccharine), anti-static-agents, lubricants (see, for example, Published PCT Patent Appl. No. WO 87/905213, the teachings of which are incorporated by reference herein), chemical stabilizers, buffers, preservatives, absorption enhancers, and other materials known to those of ordinary skill.

Sustained release coating materials (e.g., stearic acid or polymers, e.g. polyvinyl pyrrolidone, polylactic acid) may also be employed on active material or active material containing particles (see, for example, Patent Nos. US 3,634,582, GB 1,230,087, GB 1,381,872, the teachings of which are incorporated by reference herein).

Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants.

Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

Preferred unit dosage formulations are those containing an effective dose, as herein before recited, or an appropriate fraction thereof, of the active ingredient.


Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer or step or group of integers but not to the exclusion of any other integer or step or group of integers or steps.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore the Examples herein are to be construed as merely illustrative and not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is claimed is:

1. A pharmaceutical combination comprising:  
a compound according to Formula (I) hereinbelow:



(I)

wherein:

R2 and R3 are, independently, selected from the group consisting of straight or branched chain lower alkyl groups (having preferably from 1 to 6 carbon atoms), cycloalkyl groups (having from 5 to 6 carbon atoms), cycloalkyl-alkyl (having 6 to 10 carbon atoms), 2-thienyl, 2-pyridyl, phenyl, phenyl substituted with an alkyl group having not in excess of 4 carbon atoms, and phenyl substituted with an alkoxy group having not in excess of 4 carbon atoms; and

X<sup>-</sup> represents an anion associated with the positive charge of the N atom;

such that the compound is in quaternary salt form; and  
one or more other therapeutic ingredients.

2. A combination according to claim 1 wherein in the compound of formula (I) the orientation of the alkyl chain attached to the tropane ring is endo.

3. A combination according to claim 2 wherein the compound of formula (I) is selected from the group consisting of:

(3-*endo*)-3-(2,2-di-2-thienylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide;

(3-*endo*)-3-(2,2-diphenylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide;

(3-*endo*)-3-(2,2-diphenylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane 4-methylbenzenesulfonate;

(3-*endo*)-8,8-dimethyl-3-[2-phenyl-2-(2-thienyl)ethenyl]-8-azoniabicyclo[3.2.1]octane bromide;

and

(3-*endo*)-8,8-dimethyl-3-[2-phenyl-2-(2-pyridinyl)ethenyl]-8-azoniabicyclo[3.2.1]octane

bromide.

4. A combination according to claim 3 wherein in the compound of formula (I) X<sup>-</sup> is selected from the group consisting of chloride, bromide, iodide, sulfate, benzene sulfonate and toluene sulfonate.
5. A combination according to any of claims 1-4 above wherein said other therapeutic agent is selected from a  $\beta$ 2-adrenoreceptor agonist, an anti-histamine, an anti-allergic or an anti-inflammatory agent.
6. A combination according to claim 5 wherein the  $\beta$ 2-adrenoreceptor agonist is selected from salmeterol or a pharmaceutically acceptable salt thereof.
7. A combination according to claim 6 wherein the  $\beta$ 2-adrenoreceptor agonist is salmeterol xinafoate.
8. A combination according to claim 5 wherein the anti-inflammatory agent is an anti-inflammatory corticosteroid.
9. A combination according to claim 8 wherein the anti-inflammatory corticosteroid is fluticasone propionate.
10. A combination according to claim 5 wherein the anti-inflammatory agent is a PDE4 inhibitor.
11. A pharmaceutical combination according to claim 1 above wherein said other therapeutic ingredients comprise a  $\beta$ 2-adrenoreceptor agonist and an anti-inflammatory corticosteroid.
12. A pharmaceutical combination according to claim 1 wherein the  $\beta$ 2-adrenoreceptor agonist is salbutamol, salmeterol, fenoterol, terbutaline, or a salt thereof; and the anti-inflammatory corticosteroid is fluticasone propionate.
13. A combination according to claim 12 wherein the  $\beta$ 2-adrenoreceptor agonist is salmeterol xinafoate,
14. A combination according to claim 1 comprising (3-*endo*)-3-(2,2-di-2-thienylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide salmeterol xinafoate, and fluticasone propionate.
15. A pharmaceutical composition comprising a combination as defined in any of claims 1 – 14 together with one or more pharmaceutically acceptable carriers and/or excipients.

16. A method of inhibiting the binding of acetylcholine to its receptors in a mammal in need thereof comprising administering a safe and effective amount of a composition according to claim 1.

5

17. A method of treating a muscarinic acetylcholine receptor mediated disease, wherein acetylcholine binds to said receptor, comprising administering a safe and effective amount of a composition according to claim 1.

10

18. A method according to claim 17 wherein the disease is selected from the group consisting of chronic obstructive lung disease, chronic bronchitis, asthma, chronic respiratory obstruction, pulmonary fibrosis, pulmonary emphysema and allergic rhinitis.

15

19. A method according to claim 17 wherein administration is via inhalation via the mouth or nose.

20

20. A method according to claim 17 wherein administration is via a medicament dispenser selected from a reservoir dry powder inhaler, a multi-dose dry powder inhaler or a metered dose inhaler.

21. A method according to claim 17 wherein the compound is administered to a human and has a duration of action of 12 hours or more for a 1 mg dose.

25

22. A method according to claim 21 wherein the compound has a duration of action of 24 hours or more.

23. A method according to claim 22 wherein the compound has a duration of action of 36 hours or more.